

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

27 JAN 2005

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
12 February 2004 (12.02.2004)

PCT

(10) International Publication Number
WO 2004/013276 A2

(51) International Patent Classification⁷:

C12N

(71) Applicants and

(21) International Application Number:

PCT/IB2003/003681

(72) Inventors: PRASSLER, Josef [DE/DE]; Sandstr. 20, 82110 Munich (DE). STARK, Yvonne [DE/DE]; Hohangerstr. 10, 81735 Munich (DE).

(22) International Filing Date: 30 July 2003 (30.07.2003)

(81) Designated States (national): AU, CA, US.

(25) Filing Language:

English

Published:

— without international search report and to be republished upon receipt of that report

(26) Publication Language:

English

(48) Date of publication of this corrected version:

1 April 2004

(30) Priority Data:


60/399,150 30 July 2002 (30.07.2002) US

(15) Information about Correction:


see PCT Gazette No. 14/2004 of 1 April 2004, Section II

(71) Applicant (for all designated States except US): MORPHOSYS IP GMBH [DE/DE]; Lena-Christ-Str. 48, 82152 Martinsried/Planegg (DE).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



(54) Title: NOVEL TRICISTRONIC VECTORS AND USES THEREFOR



(57) Abstract: A tricistronic vector (i.e., a vector capable of expressing three exogenous genes, which are not fused together, under the control of one promoter) effectively can encode an immunoglobulin-presenting polypeptide and two immunoglobulin (Ig) polypeptides. The encoded Ig-presenting polypeptide is able to associate with at least one of the Ig polypeptides via co-expressed associating agents. A vector according to the present invention particularly is suited for phage display technology, e.g., when the Ig-presenting polypeptide is a phage coat protein and the Ig polypeptides associate to form a Fab.

WO 2004/013276 A2